



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
29.12.1999 Bulletin 1999/52

(51) Int. Cl.<sup>6</sup>: **C12N 15/81**, C12P 25/00,  
C12N 15/55, C12N 9/78,  
A23K 1/16, A23L 1/302

(21) Application number: **99107615.9**

(22) Date of filing: **16.04.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: **23.04.1998 EP 98107380**

(71) Applicant:  
**F. HOFFMANN-LA ROCHE AG**  
**4070 Basel (CH)**

(72) Inventors:  
• **Babyak, Lyubov Ya.**  
**290056 Lviv (UA)**  
• **Bacher, Adelbert**  
**85748 Garching (DE)**  
• **Boretskyy, Yuriy R.**  
**290035 Lviv (UA)**  
• **Demchyshyn, Vasyl V.**  
**290040 Lviv (UA)**

• **Eberhardt, Sabine**  
**85737 Ismaning (DE)**  
• **Fedorovych, Dariya**  
**290000 Lviv (UA)**  
• **Luettgen, Holger**  
**85748 Garching (DE)**  
• **Richter, Gerald**  
**85080 Gaimersheim (DE)**  
• **Van Loon, Adolphus**  
**4310 Rheinfelden (CH)**

(74) Representative: **Braun, Axel et al**  
**F.Hoffmann-La Roche AG**  
**Patent Department (PLP),**  
**124 Grenzacherstrasse**  
**4070 Basel (CH)**

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) **Overproduction of riboflavin in yeast**

(57) The present invention is directed to a yeast strain which has been transformed by a recombinant DNA sequence comprising a DNA sequence which upon expression in a suitable host cell encodes at least one polypeptide with riboflavin biosynthetic activity and which DNA sequence is transcriptionally linked to a promoter functional in such yeast strain, a process for the production of riboflavin characterized therein that such a yeast strain is cultured under suitable culture conditions and process for the production of a food or feed composition

**EP 0 967 287 A2**

## Description

[0001] Derivatives of riboflavin (the flavocoenzymes FMN and FAD) are universally required for redox reactions in all cellular organisms. Riboflavin (vitamin B<sub>2</sub>) is produced by all plants and by many microorganisms [Demain A.L. Riboflavin oversynthesis. Ann. Rev. Microbiol. 1972, 26, 369]. The compound is not produced in vertebrates. Riboflavin is therefore an essential nutrient for man and animals.

[0002] Riboflavin can be produced by chemical synthesis and by various fermentation procedures using strains of *Bacillus* (e.g. *Bacillus subtilis*), the ascomycetes *Ashbya gossypii* and *Eremothecium ashbyi* [Demain A.L. Riboflavin oversynthesis. Ann. Rev. Microbiol. 1972, 26, 369 and Mitsuda H, Nakajima K., Effects of 8-azaguanine on riboflavin production and on the nucleotide pools in non-growing cells of *Eremothecium ashbyi*. J. Nutr Sci Vitaminol (Tokyo) 1973; 19(3):215-227], various yeast strains such as *Candida guilliermondii*, *Candida famata* [F.W. Tanner, Jr.C. Vojnovich, J.M. Van Lanen. Riboflavin production by *Candida* species. Nature, 1945, 101 (2616):180-181] and related strains, as well as other microorganisms.

[0003] The pathway of riboflavin biosynthesis in yeast is shown in Fig. 1. The precursors for the biosynthesis of the vitamin are guanosine triphosphate (GTP) and ribulose 5-phosphate. One mol of GTP and two mol of ribulose 5-phosphate are required to biosynthetically generate one mol of riboflavin.

[0004] In the yeast, *Saccharomyces cerevisiae*, the biosynthesis of the vitamin requires at least six genes, specifically the genes *RIB1*, *RIB2*, *RIB3*, *RIB4*, *RIB5* and *RIB7* [Oltmanns O., Bacher A., Lingens F. and Zimmermann F.K. Biochemical and genetic classification of riboflavine deficient mutants of *Saccharomyces cerevisiae*. Mol. Gen. Genet. 1969, 105, 3061. In *C. guilliermondii*, the biosynthesis of riboflavin has also been shown to require the products of at least six genes, specifically the genes *RIB1*, *RIB2*, *RIB3*, *RIB4*, *RIB5* and *RIB6* (2). The enzymes specified by these *C. guilliermondii* genes and their roles in the biosynthetic pathway are summarized in Fig. 1. In contrast to the situation in *B. subtilis*, the riboflavin biosynthetic genes are not clustered in the eukaryotes *S. cerevisiae* and *C. guilliermondii*.

[0005] The initial step in the biosynthetic pathway is the opening of the imidazole ring of GTP catalyzed by the enzyme, GTP cyclohydrolase II. The product of this enzyme has been reported to be 2,5-diamino-6-ribosylamino-4(3H)-pyrimidinone 5'-phosphate. This intermediate is converted to 5-amino-6-ribitylamino-2,4(1H,3H)-pyrimidinedione by a sequence of side chain reduction, ring deamination, and dephosphorylation. The hypothetical enzyme involved in the dephosphorylation of 5-amino-6-ribitylamino 5'-phosphate is still unknown. The conversion of 5-amino-6-ribitylamino-2,4(1H,3H)-pyrimidinedione to 6,7-dimethyl-8-ribityllumazine by the enzyme, 6,7-dimethyl-8-ribityllumazine synthase, requires a second substrate, 3,4-dihydroxy-2-butanone 4-phosphate, which is obtained from ribulose 5-phosphate by the catalytic action of 3,4-dihydroxy-2-butanone-4-phosphate synthase. Finally, 6,7-dimethyl-8-ribityllumazine is converted to riboflavin by a dismutation reaction catalyzed by riboflavin synthase. The sequence of the *RIB1* gene directing the synthesis of GTP cyclohydrolase II, the initial enzyme of the riboflavin pathway, has been established in the yeast, *C. guilliermondii* (4).

[0006] Recombinant strains of *Bacillus subtilis* for the production of riboflavin by fermentation have been described, e.g. in EP 405 370. These strains carry the riboflavin operon under the control of a strong promoter directing the production of the cognate enzymes in large amount. The gene constructs of the riboflavin operon under the control of a strong promoter can be present at one or several different locations on the *B. subtilis* chromosome. The incorporation of an additional gene of the riboflavin pathway under the control of a strong promoter at a separate locus on the *B. subtilis* chromosome has also been shown to increase the yield of riboflavin obtained by fermentation, see EP 821 063.

[0007] Whereas the production of riboflavin by strains of yeasts such as *C. guilliermondii* has been reported, recombinant DNA technology has not been applied for the overexpression of riboflavin biosynthetic genes in *C. guilliermondii* or in related flavinogenic yeasts so far.

[0008] It is therefore an object of the present invention to provide recombinant means which should allow the production of yeast strains which overproduce riboflavin. More specifically it is an object of the present invention to provide a yeast strain which has been transformed by a recombinant DNA sequence comprising a DNA sequence which upon expression in a suitable host cell encodes at least one polypeptide with riboflavin biosynthetic activity and which DNA sequence is transcriptionally linked to a promoter functional in such yeast strain and even more specifically such a yeast strain which belongs to the groups of flavinogenic yeasts which overproduce riboflavin under conditions of iron starvation-like a yeast strain which is selected from the following group: *Schwanniomyces*, preferably *Schwanniomyces occidentalis*, *Debaryomyces*, preferably *Debaryomyces hansenii*, *Torulopsis*, preferably *Torulopsis candida*, or, especially *Candida*, preferably *Candida guilliermondii* or *Candida famata* [Logvinenko et al., Ukrainskii Biokhimicheskii Zhurnal 61(1), 28-32, 1989; Logvinenko et al., Mikrobiologiya 57(2), 181-186, 1988 and Nakase and Suzuki, Journal of General and Appl. Mikrobiology 31(1), 49-70 (1985)]. It is furthermore an object of the present invention to provide such yeast strains wherein the polypeptide encoding DNA sequence is from yeast, preferably flavinogenic yeasts which overproduce riboflavin under conditions of iron starvation, more preferably *Candida*, e.g. *Candida guilliermondii* or *Candida famata*.

[0009] It is also an object of the present invention to provide such yeast strains wherein the polypeptide encoding DNA

bacteria

yeast strain  
into yeast

sequence encodes a protein with GTP cyclohydrolase II activity and is selected from the following DNA sequences:

a) the DNA sequence as shown in Fig. 5 or its complementary strand;

b) DNA sequences which hybridize under standard conditions to the protein coding regions of the DNA sequences defined in (a) or fragments thereof; and

c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) and (b).

[0010] It is furthermore an object of the present invention to provide such yeast strains wherein the promotor is the TEF *S. cerevisiae* promotor.

[0011] It is also an object of the present invention to provide a process for the production of riboflavin characterized therein that a yeast strain as described above is cultured under suitable culture conditions and the riboflavin produced is isolated from the medium or the yeast strain by methods known to the man skilled in the art, and a process for the production of a food or feed composition characterized therein that riboflavin as obtained by such process is mixed with one or more suitable food or feed ingredients by a process known to the man skilled in the art.

[0012] All *C. guilliermondii* strains used in the practice of the present invention are derivatives of the *C. guilliermondii* strain obtained from the American Type Culture Collection (ATCC) under accession No. ATCC 9058 (1): *Candida guilliermondii* (ATCC 9058) has been redeposited as a Budapest Treaty deposit on April 1, 1998 and has been assigned accession No. ATCC 74437. *Candida guilliermondii* is a representative of yeast species which overproduce riboflavin (vitamin B2) under conditions of iron starvation. The group includes also *Schwanniomyces occidentalis*, (or called *Debaryomyces occidentalis*) *Debaryomyces cloeckeri*, *Torulopsis candida* and *Candida famata*. The latter species is used for industrial production of riboflavin. Regarding the taxonomic assignments of yeast species a man skilled in the art knows that these assignments are handled variably by different authors, for example: *Candida famata* - *Debaryomyces hansenii* - *Torulaspora hansenii*, *Candida guilliermondii* - *Pichia guilliermondii* - *Yamadazyma guilliermondii* are used as synonyms. A man skilled in the art knows that microorganisms which can be used for the practice of the present invention, either as host cells or source for the isolation of DNA sequences, are available from depository authorities, e.g. the American Type Culture Collection (ATCC), the Centraalbureau voor Schimmelcultures (CBS) or the Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH (DSMZ) or any other depository authority as listed in the Journal "Industrial Property" [(1991) 1, pages 29-40].

[0013] DNA sequences useful for the practice of the present invention and encoding a polypeptide with riboflavin biosynthetic activity can be obtained from any microorganism known to produce riboflavin (see above) in form of e.g., genomic or c-DNA sequences by methods known to the man skilled in the art or by using the wellknown PCR-Technology. The principles of the polymerase chain reaction (PCR) method are outlined e.g. by White et al., Trends in Genetics, 5, 185-189 (1989), whereas improved methods are described, e.g. in Innis et al. [PCR Protocols: A guide to Methods and Applications, Academic Press, Inc. (1990)].

[0014] The sequence information needed for the design of the PCR-primers can be obtained from any sequence data base, for example like Genbank (Intelligenetics, California, USA), European Bioinformatics Institute (Hinxton Hall, Cambridge, GB), NBRF (Georgetown University, Medical Centre, Washington DC, USA) and Vecbase (University of Wisconsin, Biotechnology Centre, Madison, Wisconsin, USA).

[0015] Once such DNA sequences have been obtained they can be expressed in any desirable host and the riboflavin biosynthetic activity of the encoded polypeptide can be determined by any assay known to the man skilled in the art and described, e.g. in Bacher A., G. Richter, H. Ritz, S. Eberhardt, M. Fisher and C. Krieger, Biosynthesis of riboflavin: GTP cyclohydrolase II, deaminase, and reductase. Methods in enzymology 1997; 280: 382-389; K. Kis, R. Volk and A. Bacher, Biosynthesis of riboflavin. Studies on the reaction mechanism of 6,7-dimethyl-8-ribityllumazine synthase. Biochemistry 1995, 34, 2883-2892; Logvinenko EM, Shavlovskii GM, Zakal'skii AE, Kontorovskaia Niu. Properties of 2,5-diamino-4-oxy-6-ribosylaminopyrimidine-5'-phosphate reductase, a enzyme of the second stage of flavinogenesis in *Pichia guilliermondii* yeast Ukr Biokhim Zh 1989 Jul; 61(4): 47-54; G. Richter, M. Fischer, C. Krieger, S. Eberhardt, H. Lüttgen, I. Gerstenschläger and A. Bacher. Biosynthesis of riboflavin. Characterization of the bifunctional deaminase/reductase of *Escherichia coli* and *Bacillus subtilis*. J. Bacteriol. 1997, 179, 2022-2028; K. Ritsert, D. Turk, R. Huber, R. Ladenstein, K. Schmidt-Bäse and A. Bacher. Studies on the lumazine synthase/riboflavin synthase complex of *Bacillus subtilis*. Crystal structure analysis of reconstituted icosahedral  $\beta$  subunit capsid at 2.4 Å resolution. J. Mol. Biol. 1995, 253, 151-167.

[0016] The DNA sequences used for the practice of the present invention comprise at least one DNA sequence which encodes a polypeptide with riboflavin biosynthetic activity. It is however, understood by the man skilled in the art that also more than one, e.g. all enzymes of the riboflavin biosynthetic pathway can be encoded by such DNA sequences and one or more of these enzymes can be encoded by DNA sequences of different species origin or can be of partial or total

synthetic origin as long as they show at least one desired riboflavin biosynthetic activity. One example of such a DNA sequence is given in Fig. 5 coding for a GTP cyclohydrolase II. However, DNA sequences which hybridize under standard conditions to this DNA sequence and encode such a GTP cyclohydrolase are also useful for the practice of the present invention.

- 5 [0017] "Standard conditions" for hybridization in this context are conditions which are generally used by a man skilled in the art to detect specific hybridization signals and which are described, e.g. by Sambrook et al., "Molecular Cloning" second edition, Cold Spring Harbor Laboratory Press 1989, New York, or preferably so called stringent hybridization and non-stringent washing conditions or more preferably so called stringent hybridization and stringent washing conditions a man skilled in the art is familiar with and which are described, e.g. in Sambrook et al. (s.a.). "Fragment of the
- 10 DNA sequences" means in this context a fragment which codes for a polypeptide still having the enzymatic activity as specified above.

- [0018] For the overexpression of the proteins encoded by the DNA sequences of the present invention these sequences can be linked to promoters which are functional in the desired yeast and are, e.g. the *S.cerevisiae* TEF-promotor (see Example 2) or the *pho5*-promotor [Vogel et al., Mol. Cell. Biol., 2050-2057 (1989); Rudolf and Hinnen, Proc. Natl. Acad. Sci. 84, 1340-1344 (1987)] or the *gap*-promotor or the *aox1*-promotor [Koutz et al., Yeast 5, 167-177 (1989); Sreekrishna et al., J. Basic Microbiol. 28, 265-278 (1988)] or the FMD promotor [Hollenberg et al., EP 299108] or MOS-promotor [Ledeboer et al., Nucleic Acids Res. 13, 3063-3082 (1985)].

- [0019] The DNA sequences useful for the practice of the present invention can also comprise so called "ARS" elements (autonomously replicating sequence) as described, e.g. in Example 1.

TEF promoter  
pho5 promoter  
gap promoter  
aox1 promoter  
fmd promoter  
ARS  
elements

### Examples

- [0020] If not specifically indicated or referred to by references standard procedures have been used as described, e.g. in Sambrook et al. "Molecular Cloning", (s.a.) and Cregg, J.M., K.J. Barriner, A.Y.Hessler, and K.R. Madden (1985).
- 25 *Pichia pastoris* as a host system for transformations. Mol. Cell. Biol. 5, 3376-3385.

### Example 1

#### Autonomous replication of plasmid p19R1 in *C. guilliermondii*

- 30 [0021] The plasmid pFR1 carrying the *RIB1* gene of *C. guilliermondii* has been described (Zakalsky et al. Genetika 26, 614-620, 1990). In order to subclone the *RIB1* gene, plasmid pFR1 was digested with the restriction nuclease *Sall*. The resulting fragments were cloned into the *Sall* site of the pUC19 vector. The ligation mixture was transformed into the *E. coli* mutant strain BSV821 carrying a mutation of the *ribA* gene conducting to riboflavin deficiency. Colonies growing
- 35 in the absence of riboflavin were isolated and were shown to contain a plasmid p19R1.

- [0022] The plasmid p19R1 was sequenced and was shown to contain a 2.18 kb fragment of *C. guilliermondii* DNA in the *Sall* site of the pUC19 vector. The sequence of this insert is shown in Fig. 2.

- [0023] The DNA sequence shown in Fig. 2 carries the *RIB1* gene of *C. guilliermondii*. The plasmid transforms *C. guilliermondii* mutant defective in the *RIB1* gene to riboflavin prototrophy and can replicate autonomously in this yeast species. The replication was shown to be due to the presence of an autonomously replicating sequence (ARS) comprising approximately base pairs 1542 to 1755 in Fig. 2 and extending into the structural gene *RIB1*.

rib1 →  
p19R1

### Example 2

- 45 **Construction of a plasmid for hyperexpression of the *RIB1* gene of *C. guilliermondii***

[0024] The *TEF* gene of the yeast, *S. cerevisiae*, specifies the translation elongation factor 1-alpha. This gene is known to be transcribed in *S. cerevisiae* with high efficiency.

- 50 [0025] A DNA fragment carrying the *S. cerevisiae* *TEF* promoter and the 5' part of the the *RIB1* gene of *C. guilliermondii* was obtained by PCR amplification. Initially, a DNA sequence located upstream from the 5' end of the *S. cerevisiae* *TEF* gene was amplified by PCR with primers ShBle\_V and TEF1\_H using chromosomal DNA of *S. cerevisiae* as a template. The amplified DNA fragment (subsequently designated *TEF* promoter) comprises bp 15 984 to 16 344 of the sequence listed under EMBL accession number gb/U51033/YSCP9513.

- [0026] Independently, a DNA fragment comprising the 5'-terminal part of *C. guilliermondii* GTP cyclohydrolase II structural gene was obtained by PCR with primers PGgtpCY\_V and PGgtpCY\_nco using p19R1 plasmid as a template. The amplified DNA fragment (subsequently designated 5'GTPcII) comprises bp 460 to 1145 of the sequence shown in Fig. 2.

[0027] The DNA amplicates obtained in the two PCR reactions described above comprising parts of the *TEF* gene

of *S. cerevisiae* and of the 5'GTPcII were mixed, and a third PCR amplification was performed using the primers ShBle\_V and PGgtpCY\_nco. This reaction yielded a DNA fragment which contains the complete *TEF* promoter and the 5' part of the *C. guilliermondii* *RIB1* gene.

[0028] All primers used in these PCRs are shown in Table 2. The sequence of the final amplificate is shown in Fig. 3.

[0029] The amplificate contains a cutting site for the restriction nucleases *SphI* which had been introduced via the primer ShBle\_V. The amplificate also contains an *MscII* site which is a feature of the *RIB1* gene. The amplificate was digested with *SphI* and *MscI*.

[0030] The plasmid p19R1 (whose construction is described above) was also digested with the same enzymes, and the PCR amplificate was ligated into the digested plasmid.

[0031] The ligation mixture was transformed into a mutant designated Rib7 of *Escherichia coli* carrying a mutation of the *ribA* gene which specifies GTP cyclohydrolase II. Transformation was performed by electroporation according to the protocol of Invitrogen (5).

[0032] The *E. coli* cells were plated on Luria-Broth plates supplemented with ampicillin (100 mg/ml) which did not contain riboflavin. Colonies growing on this medium were isolated and were shown to contain a plasmid designated pTC2.

[0033] The sequence of the insert of the plasmid pTC2 is shown in Fig. 3.

[0034] Plasmid pTC2 was digested with *XhoI* and *Sall* restriction endonucleases yielding 4.4 kb and 0.5 kb fragments. The 4.4 kb fragment was circularized with T4 DNA ligase. The ligation mixture was transformed into the Rib7 mutant of *E. coli* carrying the RibA mutation. Transformation was achieved by electroporation. The cells were plated on Luria-Broth plates containing ampicillin and no riboflavin (see above). Colonies growing on these plates were isolated and were shown to contain a plasmid pTCdXS2. The procedure had resulted in the removal of 0.5 kb base pairs from the plasmid pTC2.

[0035] The sequence of the insert of the plasmid pTCdXS2 is shown in Fig. 4. The open reading frame of the *RIB1* gene of *C. guilliermondii* is indicated. The DNA segment representing the promoter of the *TEF* gene of *S. cerevisiae* is also shown.

### Example 3

#### Construction of a recombinant *C. guilliermondii* strains

[0036] The riboflavin deficient mutant rh-21 with an apparent defect of the *RIB1* gene specifying GTP cyclohydrolase II (2) has been obtained after chemical mutagenesis of the L2 strain (2) which was previously obtained from the ATCC 9058 *C. guilliermondii* strain.

[0037] The plasmid pTCdXS2 was transformed into the *RIB1* mutant strain rh-21 of *C. guilliermondii* by the LiCl procedure, respectively (6). The cells were plated on YPD medium without added riboflavin. Colonies growing without riboflavin were isolated. They were monitored for GTP cyclohydrolase II activity and for riboflavin production as described below.

[0038] The prototrophic strains were monitored for the presence of DNA segments introduced with the plasmid by PCR analysis. PCR was performed using the primers ShBle\_V and PGgtpCY\_nco and boiled *C. guilliermondii* recombinant strains cells as template. Primer ShBle\_V is complementary to the *TEF* promoter and primer PGgtpCY\_nco is complementary to the *RIB1* structural gene. Amplificates of the expected length (1175 base pairs) were obtained from all transformants isolated. The amplificate obtained from strain XS-3 was isolated and was sequenced by the fluorescent dideoxy terminator method. The sequence is shown in Fig. 5. This sequence is identical with base pairs 1 to 1168 of the insert of plasmid pTCdXS2.

[0039] The recombinant transformants were genetically stable. Specifically, they did not segregate riboflavin deficient subclones.

### Example 4

#### GTP cyclohydrolase activity in recombinant *C. guilliermondii* strains

[0040] The level of GTP cyclohydrolase II activity in the recombinant strains described above was determined as follows.

[0041] The recombinant *C. guilliermondii* cells were grown aerobically in synthetic Burkholder medium supplemented with trace elements (Science 101, 180, 1945) but without asparagine, during 2-3 days at 30°C. *C. guilliermondii* L2 strain (wild type) served as a control in these experiments.

[0042] Cells from exponential growth phase were harvested by centrifugation (5000 g, 15 min), washed twice with 20 mM Tris HCl, pH 8.2, containing 1 mM DTT and 1 mM MgCl<sub>2</sub>. Cells were stored at -20°C. Frozen cells mass (1-3 g) was thawed in 3-9 ml of washing buffer. Cells were disrupted by agitation with glass beads (d = 0.8 mm). After centrifugation,

rib  
rib1 (GTPase)  
p19R1  
pg. rib7  
p.g  
transformed by invivo  
pg rib1  
E. coli  
rib1 pg.

ribA  
rh-21 pg  
mutant  
rib1  
GTP cyclo  
mutant  
ribe  
prototrophic  
pg rib1 + pg

ribA

cell extract was dialyzed overnight against 100 volumes of washing buffer. Protein concentration was measured by the Lowry method.

[0043] Reaction mixtures for GTP cyclohydrolase assays contained 20 mM Tris HCl, pH 8.2, 3 mM DTT, 2 mM MgCl<sub>2</sub>, 1 mM GTP, and protein (protein concentration, 1-3 mg/ml, total volume, 4 ml). They were incubated at 37°C for 20 min in the dark.

[0044] After incubation, 2 ml aliquots were removed, and 2,3-butanedione was added to a final concentration of 0.5 mg/ml. The mixtures were incubated at 95°C for 30 min. Blank values were processed in the same way but without added diacetyl.

[0045] Differences in specific fluorescence of both types of aliquots were determined and were used to calculate the concentrations of 6,7-dimethylpteridin and the activity of GTP cyclohydrolase II. Results are shown in Table 3.

[0046] Strain L2 from which the mutant rh-21 had been derived was used as a control. The enzyme activity in strain L2 was 2,9 nmol mg<sup>-1</sup> h<sup>-1</sup>. No enzyme activity was found in the riboflavin deficient recipient strain rh-21 carrying a mutation of the *RIB1* gene. The recombinant strains obtained by transformation with plasmid pTCdXS2 showed enzyme levels between 6,5 and 13,4 nmol mg<sup>-1</sup> h<sup>-1</sup>. Thus, the enzyme level in recombinant strains was 2.3 - 4.6-fold higher as compared with the *C. guilliermondii* strain L2.

[0047] The activity of riboflavin synthase was also measured in the recombinant strains. The activity of riboflavin synthase was not affected by the transformation with the plasmids p19RI, pTC2, pTCdXS2. All strains analyzed had riboflavin synthase activities in the range of 20 nmol mg<sup>-1</sup> h<sup>-1</sup> (Table 3).

rib1 → pg

rib 7  
pg +

#### Example 5

##### Production of riboflavin by recombinant *C. guilliermondii* strains

[0048] *C. guilliermondii* strains (wild type and recombinant strains) were grown aerobically in synthetic Burkholder medium supplemented with trace elements [Science 101, p. 180, (1945)] but without asparagine during 4 days at 30 °C [F.W. Tanner, Jr.C. Vojnovich, J.M. Van Lanen, Riboflavin production by Candida species. Nature, 1945, 101 (2616): 180-181]. The suspension was centrifuged. Riboflavin concentration was determined fluorometrically. Results are shown in Table 4.

[0049] The wild-strain L2 produced 1.2 mg riboflavin per liter under the conditions described. The recombinant strain XS-3 produced a 3-fold increased level of riboflavin (3.6 mg/l).

pichia

#### Example 6

##### Isolation of riboflavin

[0050] Two Erlenmeyer flasks (2.5 l) each containing 0.5 l of synthetic Burkholder medium containing trace elements but no asparagine were inoculated with the recombinant *C. guilliermondii* strain XS-3. The cultures were incubated with shaking at 30°C for 50 h. The solution was centrifuged. The supernatant was passed through a column of Florisil (4 ml bed volume) at a velocity of 500 ml/h. The column was washed with distilled water (7 ml). Riboflavin was eluted by a mixture of acetone/1M aqueous NH<sub>4</sub>OH. The effluent was evaporated to dryness. The yield of riboflavin was determined photometrically.

pichia

#### References

##### [0051]

1. Sibirny, A. A., Zharova, V. P., Kshanovskaya, B. V., Shavlovsky, G. M. Selection of a genetic line of yeast *Pichia guilliermondii* capable to formation of significant amount of spores. *Tsytologia i genetika* 11, 330-333, 1977 (in Russian).

2. Shavlovskyy, G. M., Sibirnyy, A. A., Kshanov'ska, B. V. Genetical classification of *C. guilliermondii* riboflavin auxotroph mutants. *Genetika* 15, 1561-1568, 1979 (in Russian).

3. Zakalsky, A. E., Zlochevsky, M. L., Stasiv, Y. Z., Logvynenko, E. M., Beburow, M. Y., Shavlovsky, G. M. Cloning of the *Pichia guilliermondii* *RIB1* gene coding for GTP cyclohydrolase II - first enzyme of flavinogenesis - in *E. coli* cells. *Genetika* 26, 614-620, 1990 (in Russian).

4. Liauta-Teglivets, O., Hasslacher, M., Boretskyy, Y., Kohlwein, S. D., Shavlovskii, G. M. Molecular cloning of the

GTP cyclohydrolase. Structural gene *RIB1* of *Pichia guilliermondii* involved in riboflavin biosynthesis. *Yeast* 11, 945-952, 1995.

5. Zero Background/Kan Cloning Kit. Version A. 151204. Instruction manual. INVITROGENE.

6. Logvinenko, E. M., Stasiv, Yu. Z., Zlochevsky, M. L., Voronovsky, A. Ya., Beburow, M. Yu., Shavlovsky, G. M. Cloning of the *RIB7* gene encoding the riboflavin synthase of the yeast *Pichia guilliermondii*. *Genetika* 29, 922-927, 1993 (in Russian).

Table 1

Enzymes and genes of the riboflavin pathway				
	Enzyme	Gene		
		<i>S. cerevisiae</i>	<i>C. guilliermondii</i>	<i>E. coli</i>
A	GTP cyclohydrolase	<i>RIB1</i>	<i>RIB1</i>	<i>ribA</i>
B	bacterial deaminase			<i>ribD</i>
C	yeast reductase	<i>RIB7</i>	<i>RIB2</i>	
D	yeast deaminase	<i>RIB2</i>	<i>RIB3</i>	
E	bacterial reductase			<i>ribD</i>
F	unknown phosphatase			
G	lumazine synthase	<i>RIB4</i>	<i>RIB5</i>	<i>ribE</i>
H	riboflavin synthase	<i>RIB5</i>	<i>RIB7</i>	<i>ribC</i>
I	3,4-dihydroxy-2-butanon 4-phosphate synthase	<i>RIB3</i>	<i>RIB6</i>	<i>ribB</i>

Table 2

Nucleotide sequences of the primers used.		
N	Primer	Sequence (5' - 3')
1	ShBle_V	GGGCATGCAATTCGAGCTCGGTACCCG
2	TEB1_H	CGACTCACTATAGGAGGAAGCTTGGCGC
3	PGgtpCY_V	AGGAGGAAGCTTGGCGCTATGGCATCGAAGG
4	PGgtpCY_nco	GCTGGTCGGTTAATGGGTGAAGCTGGG

Table 3

Activity of GTP cyclohydrolase II and riboflavin synthase in <i>C. guilliermondii</i> recombinant strains (time of growth: 40-48 h).				
N	Strain	Riboflavin synthase activity nmol mg <sup>-1</sup> h <sup>-1</sup>	GTP cyclohydrolase II activity nmol mg <sup>-1</sup> h <sup>-1</sup>	Ratio*
1	L2 (wild type)	21.6	2.88	1.00
2	R1-1	n.d.	10.08	3.50
3	R1-2	n.d.	4.20	1.46
4	R1-3	20.4	8.76	3.04
5	R1-4	19.8	7.80	2.70
6	R1-5	21.6	7.80	2.70
7	TC-1	20.4	9.60	3.33
8	TC-2	n.d.	8.40	2.92
9	TC-3	n.d.	7.56	2.63
10	XS-1	22.8	13.38	4.60
11	XS-2	n.d.	12.60	4.37
12	XS-3	n.d.	6.60	2.29

n.d. not determined.

\* GTP cyclohydrolase activity of recombinant strain divided by GTP cyclohydrolase activity of strain L2

Table 4

Riboflavin production by recombinant <i>C. guilliermondii</i> strains (time of growth: 110 h. incubation temperature: 30 °C).			
N	Strain	Riboflavin production [mg/l]	Relative riboflavin production
1	L2 (wild type)	1.2	1.0
4	R1-3	1.4	1.2
5	R1-4	3.6	3.0
6	R1-5	3.0	2.5
7	TC-1	1.4	1.2
9	TC-3	1.3	1.0
12	XS-3	3.6	3.0
13	XS-4	2.0	1.7
14	XS-5	2.3	1.9



## SEQUENCE LISTING

5

&lt;110&gt; F. Hoffmann-La Roche AG

&lt;120&gt; Overproduction of riboflavin in yeast

10

&lt;130&gt; Case 20071 EP1

&lt;140&gt; EP-A 99.107615.8

&lt;141&gt; 1999-04-16

15

&lt;150&gt; 98.107380.2

&lt;151&gt; 1998-04-23

&lt;160&gt; 12

20

&lt;170&gt; PADAT Sequenzmodul, Version 1.0

25

30

35

40

45

50

55

5 <210> 1  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer  
10 <400> 1

gggcatgcaa ttcgagctcg gtacccg 27

15 <210> 2  
<211> 28  
<212> DNA  
<213> artificial sequence

20 <220>  
<223> primer  
<400> 2

25 cgactcacta taggaggaag cttggcgc 28

30 <210> 3  
<211> 31  
<212> DNA  
<213> artificial sequence

35 <220>  
<223> Primer  
<400> 3

40 aggaggaagc ttggcgctat ggcacggaag g 31

45 <210> 4  
<211> 31  
<212> DNA  
<213> artificial sequence

<220>  
<223> Primer  
50 <400> 4

aggaggaagc ttggcgctat ggcacggaag g 31

55

<210> 5  
 <211> 2181  
 <212> DNA  
 5 <213> *Candida guilliermondii*

<220>  
 <223> RIB 1 Gene

10 <400> 5

	gtcgactttc actccgaagg taggtgcggc tggaagacgt cgtcccaagt cgtatgcgtt	60
15	agctgagagc gacggaaacg aaagtgatga agattacatg ctggaataat ccatagctag	120
	tgtacttgot aatacaaccg gtaaagctag ccaattgcag cgttattcac caccgccgtg	180
	gatcgggtta gtcacgtgaa ctggccgttg ggtcctgcac gtcgcttcat tattcatata	240
20	ttagtgagag tcttcctata tcagtcagca gacgtatcgg ttgatttcag gtcaaaaaga	300
	gaaaagggtg tcttacaaaa gcgaaatagc tgatacattt ttactcacag cagcatcata	360
	tttgtggaac ctttaactt gacttttcat ttcaagcaag ttattttgaa attcaaatca	420
25	tttggaatc aaaaaagaac atctaagttc tgaaaaattg tacgaacaac gctatggcat	480
	cgaaggacat agtacatccg caaccagagc gccggcacgg gtoggaaact cacgaattta	540
	ccatgcctct cttatctect acattgacac catcccatat tccatcgcaa acgcctcaaa	600
30	ttcotccgga agtgccagca gaagtacggg atcgcttgcc ccttctgaa acgttgccgtg	660
	tggtgaaatg catggcgaga gctcgtatac cgaccaactca ggggcgggag atattttctc	720
35	atttgtacga gaataacgtt gacaataaag agcatttggc tattgttttt ggggaagatg	780
	tgcggtcgaa aacgctctat cagaacgctc ccaatgagac ccagcaagat agaattgactc	840
	gtggtgctta tgtgggcaga ttgtttcctg gaagaaccga ggcagactat gacagtgagt	900
40	ctaatttgag attgaatttc gatgaaaatg gccaaattat cagagatccg agtaccacct	960
	gtagtggatg gccattttg gcccgatttc attcggaatg ttatacgggg gaaaccgcat	1020
	ggagtgcctg ttgcgattgt ggagaacaat tcgatgaagc tggtcgggta atgggtgaag	1080
45	ctgggcacgg gtgtatcgtg taccttcgtc aggaaggctg tggaaattgga cttggggaaa	1140
	agttgaaggc ttataatttg caagacttgg gagcggatag cgtccaggcc aatttgatgt	1200
50	tacgacatcc tgctgatgcg agatcttttt cgtcgcctac agccatactc ttggaacttg	1260
	ggctcaacga gatcaagttg ttgaccaaca atcccgataa aattgctgca gttgagggaa	1320

55

	gaaacagaga ggtcaaggta gtggaacggg tgcctatggt gccgttggca tggagaagtg	1380
5	agaatggaat caagtcaaaa gagatagagg gctacttgag tgctaagatt gaaaggatgg	1440
	ggcacttgct tgaaaagcca ctcaagatat gatagaagag atgaagttaa ggacttaaga	1500
	aataaatgat gaattaaatg acgcaaattgt cactactcga ttagagaaat agctataatg	1560
10	aagaattttg catttcgcaa aatttaagat aaatgcaaaa attgcaaatt acgaaatatg	1620
	catatgatac aagacaagaa aagactacta aaagtctctc gagaagaata ctgggtaacc	1680
15	ttcatctctt gattatgcac tggggctatt catatgcaga ttgcacgcc gaggtgcagc	1740
	gtttaggcgc ggtcaacgg aagccaacgg ccgccacaaa ttgtccggaa agtcgccgaa	1800
	actgatccac tggtaaccaca gcccataag aacccctttt aatattaaaa accgttcttc	1860
20	agccactttt gatcacattg ttgcagccg ccggttgctg ccatccaaac accacgcgtc	1920
	ccccgcacct ttacggtgc cactgcatt ggaatttgca taaaacagcc tcacgaagtg	1980
	gattaatttt tagagcactc aagtcacat gctgcaatct ctgcatcatg aatgaotcc	2040
25	cgttgataca gggaaactcag accgcaagcg gcgaagagtc acaagagcgt gtgatgtgtg	2100
	tcgactctag agatccccgg gtaccgagct cgaattcact ggcogtcgtt ttacaacgtc	2160
30	gtgactggga aaacctggc g	2181

35

40

45

50

55

<210> 6  
 <211> 332  
 <212> PRT  
 <213> Candida guilliermondii

<400> 6

10	Met	Ala	Ser	Lys	Asp	Ile	Val	His	Pro	Gln	Pro	Glu	Arg	Arg	His	Gly
	1				5					10					15	
	Ser	Glu	Thr	His	Glu	Phe	Thr	Met	Pro	Leu	Leu	Ser	Pro	Thr	Leu	Thr
				20					25					30		
	Pro	Ser	His	Ile	Pro	Ser	Gln	Thr	Pro	Gln	Ile	Pro	Pro	Glu	Val	Pro
15			35					40					45			
	Ala	Glu	Val	Arg	Asp	Arg	Leu	Pro	Leu	Pro	Glu	Thr	Leu	Pro	Val	Val
		50					55					60				
	Lys	Cys	Met	Ala	Arg	Ala	Arg	Ile	Pro	Thr	Thr	Gln	Gly	Pro	Glu	Ile
		65				70					75					80
20	Phe	Leu	His	Leu	Tyr	Glu	Asn	Asn	Val	Asp	Asn	Lys	Glu	His	Leu	Ala
				85						90					95	
	Ile	Val	Phe	Gly	Glu	Asp	Val	Arg	Ser	Lys	Thr	Leu	Tyr	Gln	Lys	Arg
				100					105					110		
	Pro	Asn	Glu	Thr	Gln	Gln	Asp	Arg	Met	Thr	Arg	Gly	Ala	Tyr	Val	Gly
			115					120					125			
25	Arg	Leu	Phe	Pro	Gly	Arg	Thr	Glu	Ala	Asp	Tyr	Asp	Ser	Glu	Ser	Asn
		130					135					140				
	Leu	Arg	Leu	Asn	Phe	Asp	Glu	Asn	Gly	Gln	Leu	Ile	Arg	Asp	Pro	Ser
		145				150					155				160	
	Thr	Thr	Cys	Ser	Gly	Glu	Pro	Ile	Leu	Ala	Arg	Ile	His	Ser	Glu	Cys
30					165					170					175	
	Tyr	Thr	Gly	Glu	Thr	Ala	Trp	Ser	Ala	Arg	Cys	Asp	Cys	Gly	Glu	Gln
				180					185					190		
	Phe	Asp	Glu	Ala	Gly	Arg	Leu	Met	Gly	Glu	Ala	Gly	His	Gly	Cys	Ile
		195						200					205			
35	Val	Tyr	Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Gly	Glu	Lys	Leu
		210					215					220				
	Lys	Ala	Tyr	Asn	Leu	Gln	Asp	Leu	Gly	Ala	Asp	Thr	Val	Gln	Ala	Asn
		225				230					235					240
	Leu	Met	Leu	Arg	His	Pro	Ala	Asp	Ala	Arg	Ser	Phe	Ser	Leu	Ala	Thr
				245					250						255	
40	Ala	Ile	Leu	Leu	Asp	Leu	Gly	Leu	Asn	Glu	Ile	Lys	Leu	Leu	Thr	Asn
				260					265					270		
	Asn	Pro	Asp	Lys	Ile	Ala	Ala	Val	Glu	Gly	Arg	Asn	Arg	Glu	Val	Lys
			275					280					285			
	Val	Val	Glu	Arg	Val	Pro	Met	Val	Pro	Leu	Ala	Trp	Arg	Ser	Glu	Asn
45			290					295				300				
	Gly	Ile	Lys	Ser	Lys	Glu	Ile	Glu	Gly	Tyr	Leu	Ser	Ala	Lys	Ile	Glu
		305				310					315					320
	Arg	Met	Gly	His	Leu	Leu	Glu	Lys	Pro	Leu	Lys	Ile				
					325						330					
50																
55																

<210> 7  
 <211> 1080  
 <212> DNA  
 <213> unknown

<220>  
 <223> final amplificate

<400> 7

	caattcgagc tcggtaccgc gggatcccc acacaccata gcttcaaaat gttctactc	60
15	cttttttact ctteccagatt ttctcggact ccgcgcacgc ccgtaccact tcaaaacacc	120
	caagcacagc atactaaatt tccctctttc ttctcttagg gtgtcgtaa ttacccgta	180
	taaagggttg gaaaagaaaa aagagaccgc ctcggtttct tttcttcgtc gaaaaggca	240
20	ataaaaattt ttatcacgtt tctttttctt gaaatttttt ttttgattt tttctcttt	300
	cgatgacctc ccattgatat ttaagtcaat aaacggtctt caatttctca agtttcagtt	360
	tcatttttct tgttctatta caactttttt tacttcttgc tcattagaaa gaaagcatag	420
25	caatctaate taagggcgag ctcgaaatcg aactagtact gcagcacgtg accggcgctt	480
	agtgttgaca attaatcacc ggcatagtat atcggcatag tataatacga ctcaatatag	540
	gaggaagctt ggcgctatgg catcgaagta catagtacat ccgcaaccag agcgccggca	600
30	cgggtcggaa actcacgaat ttaccatgcc totcttatct cctacattga caccatccca	660
	tattccatcg caaacgcctc aaattcctcc ggaagtgcc gcagaagtca gggatcgctt	720
35	gcccccttct gaaacgttgc ctgtggtgaa atgcatggcg agagctcgta taccgaccac	780
	tcaggggcgc gagatatttc tccatttgta cgagaataac gttgacaata aagagcattt	840
	ggctattggt tttggggaag atgtgcggtc gaaaacgctc tatcagaaac gtcccaatga	900
40	gaccagcaa gatagaatga ctctggtggtc ttatgtgggc agattgtttc ctggaagaac	960
	cgaggcagac tatgacagtg agtctaattt gagattgaat ttcatgaaa atggccaact	1020
45	tatcagagat ccgagtacca cctgtagtgg tgagcccatt ttggcccgta ttcattcgga	1080

<210> 8  
 <211> 175  
 5 <212> PRT  
 <213> unknown

<400> 8

10	Met	Ala	Ser	Lys	Tyr	Ile	Val	His	Pro	Gln	Pro	Glu	Arg	Arg	His	Gly
	1				5					10					15	
	Ser	Glu	Thr	His	Glu	Phe	Thr	Met	Pro	Leu	Leu	Ser	Pro	Thr	Leu	Thr
				20					25					30		
	Pro	Ser	His	Ile	Pro	Ser	Gln	Thr	Pro	Gln	Ile	Pro	Pro	Glu	Val	Pro
15			35					40					45			
	Ala	Glu	Val	Arg	Asp	Arg	Leu	Pro	Leu	Pro	Glu	Thr	Leu	Pro	Val	Val
		50					55					60				
	Lys	Cys	Met	Ala	Arg	Ala	Arg	Ile	Pro	Thr	Thr	Gln	Gly	Pro	Glu	Ile
	65				70						75				80	
20	Phe	Leu	His	Leu	Tyr	Glu	Asn	Asn	Val	Asp	Asn	Lys	Glu	His	Leu	Ala
				85						90					95	
	Ile	Val	Phe	Gly	Glu	Asp	Val	Arg	Ser	Lys	Thr	Leu	Tyr	Gln	Lys	Arg
			100						105					110		
	Pro	Asn	Glu	Thr	Gln	Gln	Asp	Arg	Met	Thr	Arg	Gly	Ala	Tyr	Val	Gly
		115					120						125			
25	Arg	Leu	Phe	Pro	Gly	Arg	Thr	Glu	Ala	Asp	Tyr	Asp	Ser	Glu	Ser	Asn
		130				135						140				
	Leu	Arg	Leu	Asn	Phe	Asp	Glu	Asn	Gly	Gln	Leu	Ile	Arg	Asp	Pro	Ser
	145				150					155					160	
	Thr	Thr	Cys	Ser	Gly	Glu	Pro	Ile	Leu	Ala	Arg	Ile	His	Ser	Glu	
30					165					170					175	

35

40

45

50

55

<210> 9  
 <211> 1742  
 <212> DNA  
 <213> Candida guilliermondii

<220>  
 <223> RIB 1 Gene

<400> 9

	caattcgagc tcggtacccg gggatccccc acacaccata gottcaaaat gtttotactc	60
15	cttttttact cttccagatt ttctcggact ccgcgcacgc cogtaccact tcaaaacacc	120
	caagcacagc atactaaatt tccctctttc ttctctagg gtgtcgtaa ttaccogtac	180
	taaaggtttg gaaaagaaaa aagagaccgc ctcgtttctt tttcttcgtc gaaaaggca	240
20	ataaaaattt ttatcacggt tctttttctt gaaatttttt tttttgattt ttttctcttt	300
	cgatgacctc ccattgatat ttaagtcaat aaacggtctt caatttctca agtttcagtt	360
	tcattttttct tgtcttatta caactttttt tactttctgc tcattagaaa gaaagcatag	420
25	caatctaate taaggcgag ctcgaaatcg aactagtact gcagcacgtg accggcgctt	480
	agtgttgaca attaatcacc ggcatagtat atcggcatag tataatacga ctactatag	540
	gaggaagctt ggcgtatagg catcgaagta catagtacat ccgcaaccag agcgccggca	600
30	cggttcggaa actcacgaat ttaccatgcc tctcttatct cctacattga caccatccca	660
	tattccatcg caaacgcctc aaattcctcc ggaagtgcc gcagaagtca gggatcgctt	720
35	gccccttctt gaaacggtgc ctgtggtgaa atgcatggcg agagctcgta taccgaccac	780
	tcaggggccc gagatatttc tccatttgta cgagaataac gttgacaata aagagcattt	840
	ggctattgtt tttggggaag atgtgcggtc gaaaacgctc taccagaaac gtcccaatga	900
40	gaccagcaa gatagaatga ctcggtgtgc ttatgtgggc agattgttgc ctggaagaac	960
	cgaggcagac tatgacagtg agtotaattt gagattgaat ttcgatgaaa atggccaact	1020
	tatcagagat ccgagtacca cctgtagtgg tgagccatt ttggcccgta ttcattcgga	1080
45	atgtttatacg ggggaaaccg catggagtgc tcgttcgat tgtggagaac aattogatga	1140
	agctggtcgg ttaatgggtg aagctgggca cgggtgtatc gtgtaccttc gtcaggaagg	1200
	tcgtggaatt ggacttgggg aaaagttgaa ggcttataat ttgcaagact tgggagcggg	1260
50	taccgtccag gccaatgtga tgttacgaca tcctgctgat gcgagatctt tttcgtctgc	1320



	tacagccata ctcttgact tggggctcaa cgagatcaag ttgttgacca acaatcccg	1380
5	taaaattgct gcagttgagg gaagaaacag agaggtcaag gtagtggaac ggggtgcctat	1440
	gggtgccgttg gcatggagaa gtaagaatgg aatcaagtca aaagagatag agggctactt	1500
	gagtgcctaag attgaaagga tggggcactt gottgaaaag ccactcaaga tatgatagaa	1560
10	gagatgaagt taaggactta agaaataaat gatgaattaa atgacgcaa tgtcactact	1620
	cgattagaga aatagctata atgaagaatt ttgcatttcg caaaatttaa gataaatgca	1680
15	aaaattgcaa attacgaaat atgcatatga tacaagacaa gaaaagacta ctaaaagtct	1740
	ct	1742

20

25

30

35

40

45

50

55

<210> 10  
 <211> 332  
 <212> PRT  
 <213> unknown

<400> 10

10 Met Ala Ser Lys Tyr Ile Val His Pro Gln Pro Glu Arg Arg His Gly  
 1 5 10 15  
 Ser Glu Thr His Glu Phe Thr Met Pro Leu Leu Ser Pro Thr Leu Thr  
 20  
 Pro Ser His Ile Pro Ser Gln Thr Pro Gln Ile Pro Pro Glu Val Pro  
 35 40 45  
 15 Ala Glu Val Arg Asp Arg Leu Pro Leu Pro Glu Thr Leu Pro Val Val  
 50 55 60  
 Lys Cys Met Ala Arg Ala Arg Ile Pro Thr Thr Gln Gly Pro Glu Ile  
 65 70 75 80  
 20 Phe Leu His Leu Tyr Glu Asn Asn Val Asp Asn Lys Glu His Leu Ala  
 85 90 95  
 Ile Val Phe Gly Glu Asp Val Arg Ser Lys Thr Leu Tyr Gln Lys Arg  
 100 105 110  
 Pro Asn Glu Thr Gln Gln Asp Arg Met Thr Arg Gly Ala Tyr Val Gly  
 115 120 125  
 25 Arg Leu Phe Pro Gly Arg Thr Glu Ala Asp Tyr Asp Ser Glu Ser Asn  
 130 135 140  
 Leu Arg Leu Asn Phe Asp Glu Asn Gly Gln Leu Ile Arg Asp Pro Ser  
 145 150 155 160  
 Thr Thr Cys Ser Gly Glu Pro Ile Leu Ala Arg Ile His Ser Glu Cys  
 165 170 175  
 30 Tyr Thr Gly Glu Thr Ala Trp Ser Ala Arg Cys Asp Cys Gly Glu Gln  
 180 185 190  
 Phe Asp Glu Ala Gly Arg Leu Met Gly Glu Ala Gly His Gly Cys Ile  
 195 200 205  
 35 Val Tyr Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Gly Glu Lys Leu  
 210 215 220  
 Lys Ala Tyr Asn Leu Gln Asp Leu Gly Ala Asp Thr Val Gln Ala Asn  
 225 230 235 240  
 Leu Met Leu Arg His Pro Ala Asp Ala Arg Ser Phe Ser Leu Ala Thr  
 245 250 255  
 40 Ala Ile Leu Leu Asp Leu Gly Leu Asn Glu Ile Lys Leu Leu Thr Asn  
 260 265 270  
 Asn Pro Asp Lys Ile Ala Ala Val Glu Gly Arg Asn Arg Glu Val Lys  
 275 280 285  
 Val Val Glu Arg Val Pro Met Val Pro Leu Ala Trp Arg Ser Lys Asn  
 290 295 300  
 45 Gly Ile Lys Ser Lys Glu Ile Glu Gly Tyr Leu Ser Ala Lys Ile Glu  
 305 310 315 320  
 Arg Met Gly His Leu Leu Glu Lys Pro Leu Lys Ile  
 325 330

<210> 11  
 <211> 1175  
 5 <212> DNA  
 <213> Candida guilliermondii

<220>  
 <223> RIB 1 Gene

10 <400> 11

	gggcatgcaa ttcgagctcg gtacccgggg atccccaca caccatagct tcaaaatgtt	60
15	tctactcctt ttttactcct ccagattttc tcggactccg cgcctgcgcg taccacttca	120
	aaacacccaa gcacagcata cttaaatttc ctctttcttc ctctagggcg tcgttaatta	180
	cccgtactaa aggtttggaa aagaaaaaag agaccgcctc gtttcttttt ctctgtcgaa	240
20	aaaggcaata aaaattttta tcacgtttct tttcttgaa attttttttt ttgatttttt	300
	tctctttcga tgacctccca ttgatattta agtcaataaa cggctctcaa tttctcaagt	360
	ttcagtttca ttttcttctg totattacaa ctttttttac ttcttgctca ttagaaagaa	420
25	agcatagcaa tctaacttaa gggcgagctc gaattcgaac tagtactgca gcacgtgacc	480
	ggcgccctagt gttgacaatt aatcatcgcc atagtatatc ggcatagtat aatacgactc	540
30	actataggag gaagcttggc gctatggcat cgaagtacat agtacatccg caaccagagc	600
	gccggcacgg gtcggaaact cacgaattta ccatgcctct cttatctcct acattgacac	660
	catcccatat tccatcgcaa acgcctcaaa ttcctccgga agtgccagca gaagtcaggg	720
35	atcgcttgcc ccttcttgaa acgttgcctg tggtgaaatg catggcgaga gctcgtatac	780
	cgaccactca ggggcggag atatttctcc atttgtaaga gaataacgtt gacaataaag	840
	agcatttggc tattgttttt ggggaagatg tgcggtcgaa aacgctctat cagaaacgtc	900
40	ccaatgagac ccagcaagat agaatgactc gtggtgctta tgtgggcaga ttgtttcctg	960
	gaagaaccga ggcagactat gacagttagt ctaatttgag attgaatttc gatgaaaatg	1020
	gccaaacttat cagagatccg agtaccacct gtagtggtga gcccaatttg gcccgatctc	1080
45	attcggaatg ttatacgggg gaaaccgcat ggagtgcctg ttgcgattgt ggagaacaat	1140
	tcgatgaagc tggtcggtta atgggtgaag ctggg	1175

50

55

<210> 12  
 <211> 204  
 <212> PRT  
 <213> unknown

<400> 12

10 Met Ala Ser Lys Tyr Ile Val His Pro Gln Pro Glu Arg Arg His Gly  
 1 5 10 15  
 Ser Glu Thr His Glu Phe Thr Met Pro Leu Leu Ser Pro Thr Leu Thr  
 20 25 30  
 Pro Ser His Ile Pro Ser Gln Thr Pro Gln Ile Pro Pro Glu Val Pro  
 35 40 45  
 15 Ala Glu Val Arg Asp Arg Leu Pro Leu Pro Glu Thr Leu Pro Val Val  
 50 55 60  
 Lys Cys Met Ala Arg Ala Arg Ile Pro Thr Thr Gln Gly Pro Glu Ile  
 65 70 75 80  
 Phe Leu His Leu Tyr Glu Asn Asn Val Asp Asn Lys Glu His Leu Ala  
 85 90 95  
 20 Ile Val Phe Gly Glu Asp Val Arg Ser Lys Thr Leu Tyr Gln Lys Arg  
 100 105 110  
 Pro Asn Glu Thr Gln Gln Asp Arg Met Thr Arg Gly Ala Tyr Val Gly  
 115 120 125  
 25 Arg Leu Phe Pro Gly Arg Thr Glu Ala Asp Tyr Asp Ser Glu Ser Asn  
 130 135 140  
 Leu Arg Leu Asn Phe Asp Glu Asn Gly Gln Leu Ile Arg Asp Pro Ser  
 145 150 155 160  
 Thr Thr Cys Ser Gly Glu Pro Ile Leu Ala Arg Ile His Ser Glu Cys  
 165 170 175  
 30 Tyr Thr Gly Glu Thr Ala Trp Ser Ala Arg Cys Asp Cys Gly Glu Gln  
 180 185 190  
 Phe Asp Glu Ala Gly Arg Leu Met Gly Glu Ala Gly  
 195 200

# Claims

1. A yeast strain which has been transformed by a recombinant DNA sequence comprising a DNA sequence which upon expression in a suitable host cell encodes at least one polypeptide with riboflavin biosynthetic activity and which DNA sequence is transcriptionally linked to a promotor functional in such yeast strain.
2. A yeast strain of claim 1 which belongs to the group of flavinogenic yeasts which overproduce riboflavin under conditions of iron starvation.
3. A yeast strain of claim 2 which is selected from the following group: Schwanniomyces, preferably Schwanniomyces occidentalis, Debaryomyces, preferably Debaryomyces hansenii, Torulopsis, preferably Torulopsis candida, or Candida, preferably Candida guilliermondii or Candida famata.
4. The yeast strain of claim 3 which is Candida guilliermondii or Candida famata.
5. A yeast strain as claimed any one of claims 1 to 4 wherein the polypeptide encoding DNA sequence is from yeast, preferably flavinogenic yeasts which overproduce riboflavin under conditions of iron starvation more preferably Candida, e.g. Candida guilliermondii or Candida famata.
6. A yeast strain as claimed in any one of claims 1 to 4, wherein the polypeptide encoding DNA sequence encodes a

protein with GTP cyclohydrolase II activity and is selected from the following DNA sequences:

a) the DNA sequence as shown in Fig. 5 or its complementary strand;

5 b) DNA sequences which hybridize under standard conditions to the protein coding regions of the DNA sequences defined in (a) or fragments thereof; and

c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) and (b).

10

7. A yeast strain as claimed in any one of claims 1 to 6 wherein the promotor is the TEF *S. cerevisiae* promotor.

8. A process for the production of riboflavin characterized therein that a yeast strain as claimed in any one of claims 1 to 7 is cultured under suitable culture conditions and the riboflavin produced is isolated from the medium or the yeast strain by methods known to the man skilled in the art.

15

9. A process for the production of a food or feed composition characterized therein that riboflavin as obtained by the process of claim 8 is mixed with one or more suitable food or feed ingredients.

20

25

30

35

40

45

50

55




Fig. 1

**Fig.2.**  
**Nucleotide sequence of the p19R1 insert.**

```

      10      20      30      40      50      60
      |      |      |      |      |      |
GTCGACTTTCACCTCCGAAGGTAGGTGCGGCTGGAAGACGTCGTCCCAAGTCGTATGCGTT

      70      80      90     100     110     120
      |      |      |      |      |      |
AGCTGAGAGCGACGGAACGAAAGTGATGAAGATTACATGCTGGAATAATCCATAGCTAG

      130     140     150     160     170     180
      |      |      |      |      |      |
TGTA CTGCTAATAACAACCGGTAAAGCTAGCCAATTG CAGCGTTATTCACCACCGCCGTG

      190     200     210     220     230     240
      |      |      |      |      |      |
GATCGGGTTAGTCACGTGAACTGGCCGTTGGGTCTCTGCACGTCGCTTCATTATTCATATA

      250     260     270     280     290     300
      |      |      |      |      |      |
T TAGTGAGAGTCTTCCTATATCAGTCAGCAGACGTATCGGTTGATTTCAGGTCAAAAAGA

      310     320     330     340     350     360
      |      |      |      |      |      |
GAAAAGGTGGTCTTACAAAAGCGAAATAGCTGATACATTTTTACTCACAGCAGCATCATA

      370     380     390     400     410     420
      |      |      |      |      |      |
TTTGTGGAACCTTTAAACTTGACTTTTCATTTCAAGCAAGTTATTTTGAAATTCAAATCA

      430     440     450     460     470     480
      |      |      |      |      |      |
TTTGAAATCAAAAAAGAACATCTAAGTTCTGAAAAATTGTACGAACAACGCTATGGCAT

```

MetAlaSer

```

      490     500     510     520     530     540
      |      |      |      |      |      |
CGAAGGACATAGTACATCCGCAACCAGAGCGCCGGCACGGGTCGGAACTCACGAATTTA

```

LysAspIleValHisProGlnProGluArgArgHisGlySerGluThrHisGluPheThr

```

      550     560     570     580     590     600
      |      |      |      |      |      |
CCATGCCTCTCTTATCTCCTACATTGACACCATCCCATATTCCATCGCAAACGCCTCAAA

```

METProLeuLeuSerProThrLeuThrProSerHisIleProSerGlnThrProGlnIle

610 620 630 640 650 660  
 TTCCTCCGGAAGTGCCAGCAGAAGTCAGGGATCGCTTGCCCTTCCTGAAACGTTGCCTG  
 ProProGluValProAlaGluValArgAspArgLeuProLeuProGluThrLeuProVal  
 670 680 690 700 710 720  
 TGGTGAAATGCATGGCGAGAGCTCGTATACCGACCACTCAGGGGCCGAGATATTTCTCC  
 ValLysCysMETAlaArgAlaArgIleProThrThrGlnGlyProGluIlePheLeuHis  
 730 740 750 760 770 780  
 ATTTGTACGAGAATAACGTTGACAATAAAGAGCATTGGCTATTGTTTTGGGGAAGATG  
 LeuTyrGluAsnAsnValAspAsnLysGluHisLeuAlaIleValPheGlyGluAspVal  
 790 800 810 820 830 840  
 TGCGGTGCAAAACGCTCTATCAGAAACGTCCCAATGAGACCCAGCAAGATAGAATGACTC  
 ArgSerLysThrLeuTyrGlnLysArgProAsnGluThrGlnGlnAspArgMETThrArg  
 850 860 870 880 890 900  
 GTGGTGCTTATGTGGGCAGATTGTTTCCTGGAAGAACCGAGGCAGACTATGACAGTGAGT  
 GlyAlaTyrValGlyArgLeuPheProGlyArgThrGluAlaAspTyrAspSerGluSer  
 910 920 930 940 950 960  
 CTAATTTGAGATTGAATTTTCGATGAAAATGGCCAATTATCAGAGATCCGAGTACCACCT  
 AsnLeuArgLeuAsnPheAspGluAsnGlyGlnLeuIleArgAspProSerThrThrCys  
 970 980 990 1000 1010 1020  
 GTAGTGGTGAGCCCATTTTGCCCGTATTTCATTCGGAATGTTATACGGGGGAAACCGCAT  
 SerGlyGluProIleLeuAlaArgIleHisSerGluCysTyrThrGlyGluThrAlaTrp  
 1030 1040 1050 1060 1070 1080  
 GGAGTGCTCGTTGCGATTGTGGAGAACAATTCGATGAAGCTGGTCGGTTAATGGGTGAAG  
 SerAlaArgCysAspCysGlyGluGlnPheAspGluAlaGlyArgLeuMETGlyGluAla



1090      1100      1110      1120      1130      1140  
 |      |      |      |      |      |  
 CTGGGCACGGGTGTATCGTGTACCTTCGTCAGGAAGGTCGTGGAATTGGACTTGGGGAAA  
 GlyHisGlyCysIleValTyrLeuArgGlnGluGlyArgGlyIleGlyLeuGlyGluLys

1150      1160      1170      1180      1190      1200  
 |      |      |      |      |      |  
 AGTTGAAGGCTTATAATTGCAAGACTTGGGAGCGGATACCGTCCAGGCCAATTGATGT  
 LeuLysAlaTyrAsnLeuGlnAspLeuGlyAlaAspThrValGlnAlaAsnLeuMETLeu

1210      1220      1230      1240      1250      1260  
 |      |      |      |      |      |  
 TACGACATCCTGCTGATGCGAGATCTTTTCGCTCGCTACAGCCATACTCTTGGACTTGG  
 ArgHisProAlaAspAlaArgSerPheSerLeuAlaThrAlaIleLeuLeuAspLeuGly

1270      1280      1290      1300      1310      1320  
 |      |      |      |      |      |  
 GGCTCAACGAGATCAAGTTGTTGACCAACAATCCCGATAAAATTGCTGCAGTTGAGGGAA  
 LeuAsnGluIleLysLeuLeuThrAsnAsnProAspLysIleAlaAlaValGluGlyArg

1330      1340      1350      1360      1370      1380  
 |      |      |      |      |      |  
 GAAACAGAGAGGTCAAGGTAGTGGAACGGGTGCCTATGGTGCCGTGGCATGGAGAACTG  
 AsnArgGluValLysValValGluArgValProMETValProLeuAlaTrpArgSerGlu

1390      1400      1410      1420      1430      1440  
 |      |      |      |      |      |  
 AGAATGGAATCAAGTCAAAGAGATAGAGGGCTACTTGAGTGCTAAGATTGAAAGGATGG  
 AsnGlyIleLysSerLysGluIleGluGlyTyrLeuSerAlaLysIleGluArgMETGly

1450      1460      1470      1480      1490      1500  
 |      |      |      |      |      |  
 GGCACCTTGCTTGAAGCCACTCAAGATATGATAGAAGAGATGAAGTTAAGGACTTAAGA  
 HisLeuLeuGluLysProLeuLysIle-----

1510      1520      1530      1540      1550      1560  
 |      |      |      |      |      |  
 AATAAATGATGAATTAAATGACGCAAATGTCACCTACTCGATTAGAGAAATAGCTATAATG

1570      1580      1590      1600      1610      1620  
 |      |      |      |      |      |  
 AAGAATTTTGCATTTTCGCAAAATTTAAGATAAATGCAAAAATTGCAAATTACGAAATATG

*butsky*  
*butsky top : ARS insertion*

1630 1640 1650 1660 1670 1680  
 CATATGATACAAGACAAGAAAAGACTACTAAAA**GTCTCTCGAGAAGAATACTGGGTAACC**

1690 1700 1710 1720 1730 1740  
 TTCATCTCTTGATTATGCACTGGGGCTATT**CATATGCAGATTTCGCACGCCGAGGTGCAGC**

1750 1760 1770 1780 1790 1800  
 GTTTAGGCGCGGCTCAACGGAAGCCAACGGCCGCCACAAATTGTCCGG**AACTCGCCGAA**

1810 1820 1830 1840 1850 1860  
 ACTGATCCACTGGTAC**CCACAGCCCCATAAGAACCCCTTTAATATTAA****AAACCGTTCTTC**

1870 1880 1890 1900 1910 1920  
 AGCCACTTTTGATCACATTGTTTGCAGCCGCCCGTTGCTGCCATCCAA**ACACCACGCGTC**

1930 1940 1950 1960 1970 1980  
 CCCCACAC**CTTTTACGGTGCCCACTGCATTGGAATTTGCATAAAACAGCCTCACGAAGTG**

1990 2000 2010 2020 2030 2040  
 GATTAATTTT**AGAGCACTCAAGTCATCATCTGCAATCTCTGCATCATGAAATGACTCC**

2050 2060 2070 2080 2090 2100  
 CGTTGATACAGGGA**ACTCAGACCGCAAGCGCGAAGAGTCACAAGAGCGTGTGATGTGTG**

2110 2120 2130 2140 2150 2160  
 TCGACTCTAGAGATCCCCGG**TAC**AGCTCGAATTCAGTGGCCGTCGTTTACAA**CGTC**

2170 2180  
 GTGACTGGGAAAAC**CCCTGGCG**

Sequence of the *RIB1* gene is translated.

Sequence of ARS element is typed in bold letters.

**Fig.3**  
**Nucl otide sequence of the pTC2 insert.**

```

      10      20      30      40      50      60
      |      |      |      |      |      |
CAATTCGAGCTCGGTACCCGGGATCCCCACACACCATAGCTTCAAAATGTTTCTACTC

      70      80      90     100     110     120
      |      |      |      |      |      |
CTTTTTTACTCTTCCAGATTTTCTCGGACTCCGCGCATCGCCGTACCACTTCAAAACACC

     130     140     150     160     170     180
     |      |      |      |      |      |
CAAGCACAGCATACTAAATTTCCCTCTTTCTTCCTCTAGGGTGTCGTTAATTACCCGTAC

     190     200     210     220     230     240
     |      |      |      |      |      |
TAAAGGTTTGGAAAAAGAAAAAGAGACCGCCTCGTTTCTTTTCTTCGTCGAAAAAGGCA

     250     260     270     280     290     300
     |      |      |      |      |      |
ATAAAAATTTTATCACGTTTCTTTTCTTGAAATTTTITTTTGGATTTTTCTCTTT

     310     320     330     340     350     360
     |      |      |      |      |      |
CGATGACCTCCCATTGATATTTAAGTCAATAAACGGTCTTCAATTTCTCAAGTTTCAGTT

     370     380     390     400     410     420
     |      |      |      |      |      |
TCATTTTCTTGTCTATTACAACTTTTTTACTTCTTGCTCATTACAAAGAAAGCATAG

     430     440     450     460     470     480
     |      |      |      |      |      |
CAATCTAATCTAAGGGCGAGCTCGAATTCGAACTAGTACTGCAGCACGTGACCGGCGCCT

     490     500     510     520     530     540
     |      |      |      |      |      |
ACTGTTGACAATTAATCATCGGCATAGTATATCGGCATAGTATAATACGACTCACTATAG

     550     560     570     580     590     600
     |      |      |      |      |      |
- GAGGAAGCTTGGCGCTATGGCATCGAAGTACATAGTACATCCGCAACCAGAGCGCCGCA

```

METAlaSerLysTyrIleValHisProGlnProGluArgArgHis

610 620 630 640 650 660  
 CGGGTCGGAACTCAGCAATTTACCATGCCTCTCTTATCTCTCTACATTGACACCATCCCA  
 GlySerGluThrHisGluPheThrMETProLeuLeuSerProThrLeuThrProSerHis  
 670 680 690 700 710 720  
 TATTCCATCGCAAACGCCCTCAAATTCCTCCCGAAGTCCCAGCAGAAGTCAGGGATCGCTT  
 IleProSerGlnThrProGlnIleProProGluValProAlaGluValArgAspArgLeu  
 730 740 750 760 770 780  
 GCCCCCTTCCTGAAACGTTGCCCTGTGGTGAAATGCATGCCGACAGCTCCTATACCGACCAC  
 ProLeuProGluThrLeuProValValLysCysMETAlaArgAlaArgIleProThrThr  
 790 800 810 820 830 840  
 TCAGGGGCCGGAGATATTTCTCCATTTGTACGAGAATAACGTTGACAATAAAGAGCATT  
 GlnGlyProGluIlePheLeuHisLeuTyrGluAsnAsnValAspAsnLysGluHisLeu  
 850 860 870 880 890 900  
 CGCTATTGTTTTTGGGAAGATGTGCGGTGCGAAAACGCTCTATCAGAAACGTCCTCAATGA  
 AlaIleValPheGlyGluAspValArgSerLysThrLeuTyrGlnLysArgProAsnGlu  
 910 920 930 940 950 960  
 GACCCAGCAACATAGAATGACTCGTGGTGCTTATCTGGGCAGATTGTTTCCTGGAAGAAC  
 ThrGlnGlnAspArgMETThrArgGlyAlaTyrValGlyArgLeuPheProGlyArgThr  
 970 980 990 1000 1010 1020  
 CGAGGCAGACTATGACAGTGAGTCTAATTTGACATTGAATTTGATGAAAATGGCCAACT  
 GluAlaAspTyrAspSerGluSerAsnLeuArgLeuAsnPheAspGluAsnGlyGlnLeu  
 1030 1040 1050 1060 1070 1080  
 TATCAGAGATCCGAGTACCACCTGTAOTGGTGAGCCCATTTTGGCCCGTATTCATTCGGA  
 IleArgAspProSerThrThrCysSerGlyGluProIleLeuAlaArgIleHisSerGlu

**Fig.4**  
Nucleotide sequence of the pTCdXS-2 insert.

```

      10      20      30      40      50      60
      |      |      |      |      |      |
CAATTCGAGCTCGGTACCCGGGGATCCCCACACACCATAGCTTCAAAATGTTTCTACTC

      70      80      90     100     110     120
      |      |      |      |      |      |
CTTTTTTACTCTTCCAGATTTTCTCGGACTCCGCGCATCGCCGTACCACTTCAAAACACC

     130     140     150     160     170     180
     |      |      |      |      |      |
CAAGCACAGCATACTAAATTTCCCTCTTCTTCCTCTAGGGTGTCGTTAATTACCCGTAC

     190     200     210     220     230     240
     |      |      |      |      |      |
TAAAGGTTTGAAAAAGAAAAAGAGACCGCCTCGTTTCTTTTCTTCGTCGAAAAAGGCA

     250     260     270     280     290     300
     |      |      |      |      |      |
ATAAAAATTTTTATCACGTTTCTTTTTCTTGAAATTTTTTTTTTGATTTTTTCTCTTT

     310     320     330     340     350     360
     |      |      |      |      |      |
CGATGACCTCCCATTGATATTTAAGTCAATAAACGGTCTTCAATTCTCAAGTTTCAGTT

     370     380     390     400     410     420
     |      |      |      |      |      |
TCATTTTTCTTGTTCTATTACAACCTTTTTTACTTCTTGCTCATTAGAAAAGAAAGCATAG

     430     440     450     460     470     480
     |      |      |      |      |      |
CAATCTAATCTAAGGGCGAGCTCGAATTCGAACTAGTACTGCAGCACGTGACCGGCGCCT

     490     500     510     520     530     540
     |      |      |      |      |      |
AGTGTGACAATTAATCATCGGCATAGTATATCGGCATAGTATAATACGACTCACTATAG

     550     560     570     580     590     600
     |      |      |      |      |      |
GAGGAAGCTTGGCGCTATGGCATCGAAGTACATAGTACATCCGCAACCAGAGCGCCGGCA

```

METAlaSerLysTyrIleValHisProGlnProGluArgArgHis

610 620 630 640 650 660  
 C G G G T C G G A A A C T C A C G A A T T T A C C A T G C C T C T C T T A T C T C C T A C A T T G A C A C C A T C C C A  
 GlySerGluThrHisGluPheThrMETProLeuLeuSerProThrLeuThrProSerHis

670 680 690 700 710 720  
 T A T T C C A T C G C A A A C G C C T C A A A T T C C T C C G G A A G T G C C A G C A G A A G T C A G G G A T C G C T T  
 IleProSerGlnThrProGlnIleProProGluValProAlaGluValArgAspArgLeu

730 740 750 760 770 780  
 G C C C C T T C C T G A A A C G T T G C C T G T G G T G A A A T G C A T G G C G A G A G C T C G T A T A C C G A C C A C  
 ProLeuProGluThrLeuProValValLysCysMETAlaArgAlaArgIleProThrThr

790 800 810 820 830 840  
 T C A G G G C C G G A G A T A T T T C T C C A T T T G T A C G A G A A T A A C G T T G A C A A T A A A G A G C A T T T  
 GlnGlyProGluIlePheLeuHisLeuTyrGluAsnAsnValAspAsnLysGluHisLeu

850 860 870 880 890 900  
 G G C T A T T G T T T T T G G G G A A G A T G T G C G G T C G A A A C G C T C T A T C A G A A C G T C C C A A T G A  
 AlaIleValPheGlyGluAspValArgSerLysThrLeuTyrGlnLysArgProAsnGlu

910 920 930 940 950 960  
 G A C C C A G C A A G A T A G A A T G A C T C G T G G T G C T T A T G T G G G C A G A T T G T T C C T G G A A G A A C  
 ThrGlnGlnAspArgMETThrArgGlyAlaTyrValGlyArgLeuPheProGlyArgThr

970 980 990 1000 1010 1020  
 C G A G G C A G A C T A T G A C A G T G A G T C T A A T T T G A G A T T G A A T T C G A T G A A A A T G G C C A A C T  
 GluAlaAspTyrAspSerGluSerAsnLeuArgLeuAsnPheAspGluAsnGlyGlnLeu

1030 1040 1050 1060 1070 1080  
 T A T C A G A G A T C C G A G T A C C A C C T G T A G T G G T G A G C C C A T T T T G G C C C G T A T T C A T T C G G A  
 IleArgAspProSerThrThrCysSerGlyGluProIleLeuAlaArgIleHisSerGlu

1090 1100 1110 1120 1130 1140  
 ATGTTATACGGGGAAACCGCATGGAGTGCTCGTTGCGATTGTGGAGAACAAATTCGATGA  
 CysTyrThrGlyGluThrAlaTrpSerAlaArgCysAspCysGlyGluGlnPheAspGlu  
 1150 1160 1170 1180 1190 1200  
 AGCTGGTCGGTTAATGGGTGAAGCTGGGCACGGGTGTATCGTGTACCTTCGTCAGGAAGG  
 AlaGlyArgLeuMETGlyGluAlaGlyHisGlyCysIleValTyrLeuArgGlnGluGly  
 1210 1220 1230 1240 1250 1260  
 TCGTGGAATTGGACTTGGGGAAAAGTTGAAGGCTTATAATTTGCAAGACTTGGGAGCGGA  
 ArgGlyIleGlyLeuGlyGluLysLeuLysAlaTyrAsnLeuGlnAspLeuGlyAlaAsp  
 1270 1280 1290 1300 1310 1320  
 TACCGTCCAGGCCAATTTGATGTTACGACATCCTGCTGATGCGAGATCTTTTCGCTCGC  
 ThrValGlnAlaAsnLeuMETLeuArgHisProAlaAspAlaArgSerPheSerLeuAla  
 1330 1340 1350 1360 1370 1380  
 TACAGCCATACTCTTGGACTTGGGGCTCAACGAGATCAAGTTGTTGACCAACAATCCCGA  
 ThrAlaIleLeuLeuAspLeuGlyLeuAsnGluIleLysLeuLeuThrAsnAsnProAsp  
 1390 1400 1410 1420 1430 1440  
 TAAAATTGCTGCAGTTGAGGGAAGAAACAGAGAGGTCAAGGTAGTGGAAACGGGTGCCTAT  
 LysIleAlaAlaValGluGlyArgAsnArgGluValLysValValGluArgValProMET  
 1450 1460 1470 1480 1490 1500  
 GGTGCCGTTGGCATGGAGAAGTGAGAATGGAATCAAGTCAAAAGAGATAGAGGGCTACTT  
 ValProLeuAlaTrpArgSerGluAsnGlyIleLysSerLysGluIleGluGlyTyrLeu  
 1510 1520 1530 1540 1550 1560  
 GAGTGCTAAGATTGAAAGGATGGGGCACTTGCTTGAAAAGCCACTCAAGATATGATAGAA  
 SerAlaLysIleGluArgMETGlyHisLeuLeuGluLysProLeuLysIle-----

1570	1580	1590	1600	1610	1620
<b>GAGATGAAGTTAAGGACTTAAGAAATAAATGATGAATTAAATGACGCAAATGTC</b> ACTACT					
1630	1640	1650	1660	1670	1680
<b>CGATTAGAGAAATAGCTATAATGAAGAATTTTGCATTTTCGCAAAATTTAAGATAAATGCA</b>					
1690	1700	1710	1720	1730	1740
<b>AAAATTGCAAATTACGAAATATGCATATGATACAAGACAAGAAAAGACTACTAAAAGTCTCT</b>					

Sequence of the *TEF* promoter is underlined.  
 Sequence of the *RIB1* gene is translated.  
 Sequence of ARS element is typed in bold letters.



Fig.5

Nucleotide sequence of PCR product obtained with primers ShBle\_V and PGgtpCY\_nco using boiled cells of *P.guilliermondii* XS-3 recombinant strain.

```

      10      20      30      40      50      60
      |      |      |      |      |      |
GGGCATGCAATTCGAGCTCGGTACCCGGGGATCCCCACACACCATAGCTTCAAAATGTT

      70      80      90     100     110     120
      |      |      |      |      |      |
TCTACTCCTTTTTTACTCTTCCAGATTTTCTCGGACTCCGCGCATCGCCGTACCACTTCA

      130     140     150     160     170     180
      |      |      |      |      |      |
AAACACCCAAGCACAGCATACTAAATTTCCCTCTTCTTCTCTAGGGTGTCTGTTAATTA

      190     200     210     220     230     240
      |      |      |      |      |      |
CCCGTACTAAAGGTTTGGAAAAGAAAAAGAGACCGCCTCGTTTCTTTTCTTCGTCGAA

      250     260     270     280     290     300
      |      |      |      |      |      |
AAAGGCAATAAAAAATTTTATCACGTTTCTTTTCTTGAAATTTTTTTTTTGATTTTTT

      310     320     330     340     350     360
      |      |      |      |      |      |
TCTCTTTTCGATGACCTCCCATTGATATTTAAGTCAATAAACGGTCTTCAATTTCTCAAGT

      370     380     390     400     410     420
      |      |      |      |      |      |
TTCAGTTTCATTTTTTCTGTTCTATTACAACTTTTTTTACTTCTTGCTCATTAGAAAGAA

      430     440     450     460     470     480
      |      |      |      |      |      |
AGCATEGCAATCTAATCTAAGGGCGAGCTCGAATTCGAACTAGTACTGCAGCACGTGACC

      490     500     510     520     530     540
      |      |      |      |      |      |
GGCGCCTAGTGTGACAATTAATCATCGGCATAGTATATCGGCATAGTATAATACGACTC

      550     560     570     580     590     600
      |      |      |      |      |      |
ACTATAGGAGGAAGCTTGGCGCTATGGCATCGAAGTACATAGTACATCCGCAACCAGAGC

```

METAlaSerLysTyrIleValHisProGlnProGluArg

610        620        630        640        650        660  
 |        |        |        |        |        |  
 GCCGGCACGGGTCGGAACTCACGAATTTACCATGCCTCTCTTATCTCTACATTGACAC  
 ArgHisGlySerGluThrHisGluPheThrMETProLeuLeuSerProThrLeuThrPro

670        680        690        700        710        720  
 |        |        |        |        |        |  
 CATCCCATATTCCATCGCAAACGCCTCAAATTCCTCCGGAAGTGCCAGCAGAAGTCAGGG  
 SerHisIleProSerGlnThrProGlnIleProProGluValProAlaGluValArgAsp

730        740        750        760        770        780  
 |        |        |        |        |        |  
 ATCGCTTGCCCTTCCTGAAACGTTGCCGTGGTGAAATGCATGGCGAGAGCTCGTATAC  
 ArgLeuProLeuProGluThrLeuProValValLysCysMETAlaArgAlaArgIlePro

790        800        810        820        830        840  
 |        |        |        |        |        |  
 CGACCACTCAGGGCCGGAGATATTTCTCCATTTGTACGAGAATAACGTTGACAATAAAG  
 ThrThrGlnGlyProGluIlePheLeuHisLeuTyrGluAsnAsnValAspAsnLysGlu

850        860        870        880        890        900  
 |        |        |        |        |        |  
 AGCATTGCGCTATTGTTTTGGGGAAGATGTGCGGTCGAAAACGCTCTATCAGAAACGTC  
 HisLeuAlaIleValPheGlyGluAspValArgSerLysThrLeuTyrGlnLysArgPro

910        920        930        940        950        960  
 |        |        |        |        |        |  
 CCAATGAGACCCAGCAAGATAGAATGACTCGTGGTGCTTATGTGGGCAGATTGTTTCCTG  
 AsnGluThrGlnGlnAspArgMETThrArgGlyAlaTyrValGlyArgLeuPheProGly

970        980        990        1000        1010        1020  
 |        |        |        |        |        |  
 GAAGAACCGAGGCAGACTATGACAGTGAGTCTAATTGAGATTGAATTTTCGATGAAAATG  
 ArgThrGluAlaAspTyrAspSerGluSerAsnLeuArgLeuAsnPheAspGluAsnGly


1030        1040        1050        1060        1070        1080  
 |        |        |        |        |        |  
 GCCAACTTATCAGAGATCCGAGTACCACCTGTAGTGGTGAGCCCATTTGGCCCGTATTC  
 GlnLeuIleArgAspProSerThrThrCysSerGlyGluProIleLeuAlaArgIleHis

1090        1100        1110        1120        1130        1140  
 |        |        |        |        |        |  
 ATTCGGAATGTTATACGGGGGAAACCGCATGGAGTGCTCGTTGCGATTGTGGAGAACAAT  
 SerGluCysTyrThrGlyGluThrAlaTrpSerAlaArgCysAspCysGlyGluGlnPhe

1150            1160            1170  
 |            |            |  
 TCGATGAAGCTGGTCGGTTAATGGGTGAAGCTGGG  
 AspGluAlaGlyArgLeuMETGlyGluAlaGly

Sequence of the *TEF* promoter is underlined.  
 Partial sequence of the *RIB1* gene is translated.

Scheme of the 2-steps PCR amplification.



1. SkBle-V
2. TEF1-H
3. PGgtpcy-V
4. PGgtpcy-nc0